




Europäisches
Patentamt

European
Patent Office

Office européen
des brevets

12.08.03

Rec'd PCT/PTO 07 FEB 2003
PCT/03/50373



#3

Bescheinigung

Certificate

Attestation

Die angehefteten Unterla-
gen stimmen mit der
ursprünglich eingereichten
Fassung der auf dem näch-
sten Blatt bezeichneten
europäischen Patentanmel-
dung überein.

The attached documents
are exact copies of the
European patent application
described on the following
page, as originally filed.

Les documents fixés à
cette attestation sont
conformes à la version
initialement déposée de
la demande de brevet
européen spécifiée à la
page suivante.

Patentanmeldung Nr. Patent application No. Demande de brevet n°

02078352.8

**PRIORITY
DOCUMENT**
SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH RULE 17.1(a) OR (b)

Der Präsident des Europäischen Patentamts;
Im Auftrag

For the President of the European Patent Office

Le Président de l'Office européen des brevets
p.o.

R C van Dijk

BEST AVAILABLE COPY



Anmeldung Nr:
Application no.: 02078352.8
Demande no:

Anmeldetag:
Date of filing: 14.08.02
Date de dépôt:

Anmelder/Applicant(s)/Demandeur(s):

JANSSEN PHARMACEUTICA N.V.
Turnhoutsebaan 30
2340 Beerse
BELGIQUE

Bezeichnung der Erfindung/Title of the invention/Titre de l'invention:
(Falls die Bezeichnung der Erfindung nicht angegeben ist, siehe Beschreibung.
If no title is shown please refer to the description.
Si aucun titre n'est indiqué se référer à la description.)

Use of NF-Kappa B inhibitors for the treatment of mastitis

In Anspruch genommene Priorität(en) / Priority(ies) claimed /Priorité(s)
revendiquée(s)
Staat/Tag/Aktenzeichen/State/Date/File no./Pays/Date/Numéro de dépôt:

Internationale Patentklassifikation/International Patent Classification/
Classification internationale des brevets:

A61K31/00

Am Anmeldetag benannte Vertragstaaten/Contracting states designated at date of
filing/Etats contractants désignées lors du dépôt:

AT BE BG CH CY CZ DE DK EE ES FI FR GB GR IE IT LI LU MC NL PT SE SK TR

USE OF NF-KAPPA B INHIBITORS FOR THE TREATMENT OF MASTITIS

5 The present invention relates to the use of NF- κ B (NF-kappa B) inhibitors for the manufacture of a medicament for the treatment of mastitis.

10 Mastitis is one of the most costly diseases in dairy herds. Costs mainly arise from decreased milk production and quality, therapeutic interventions, loss of antibiotic-contaminated milk and extra labor. Mastitis, defined as an inflammatory condition of the mammary gland, is usually caused by pathogenic bacteria. The most common
15 pathogens that provoke mastitis include *Staphylococcus aureus*, *Streptococcus agalactiae*, *Streptococcus dysgalactiae*, *Streptococcus uberis*, *Arcanobacterium pyogenes* and coliforms. These microorganisms can invade the udder through the teat canal and produce inflammation of the milk-producing tissue causing the formation of scar tissue which, once formed, may cause a permanent reduction in the cow's milk
production. An infection can also alter the composition, quantity, appearance and quality of the milk.

20 Mastitis in dairy cattle is usually treated with antibiotics. However it has been found to be desirable to replace such treatment with antibiotics with treatment by non-antibiotic chemo-therapeutic drugs to avoid building up of resistance of bacterial strains. Furthermore, milk produced by dairy cattle on an antibiotic treatment has to be discarded as it contains residual antibiotics.

25 NF- κ B (nuclear factor-kappa B) is a collective name for inducible dimeric transcription factors composed of members of the Rel family of DNA-binding proteins that recognize a common sequence motif. NF- κ B is found in essentially all cell types and is involved in activation of an exceptionally large number of genes in response to infections, inflammation, and other stressful situations requiring rapid reprogramming
30 of gene expression.

Surprisingly, it has been found that compounds capable of inhibiting the activation of NF- κ B, *i.e.* NF- κ B inhibitors, can be of use in the treatment of mastitis.

35 Known NF- κ B inhibitors are e.g. :

- 15d-PGJ2 (15-deoxy- Δ .12,14-prostaglandin J2) (see *Proc. Natl. Acad. Sci. USA*, 97(9), 4844-4849 (2000));
- tepoxalin (see *Immunology Letters*, 53(2,3), 109-113 (1996));

- cycloepoxydon and its (+)- and (-)- stereoisomers (see *J. Am. Chem. Soc.*, 123(45), 11308-11309 (2001));
- sodium diethyldithiocarbamate (Na-DDTC) (see *Brain Research Bulletin*, 55/3, 375-386 (2001));
- 5 - gliotoxin, MG-132 (N-[(phenylmethoxy)carbonyl]-L-leucyl-N-[(1S)-1-formyl-3-methylbutyl]-L-Leucinamide), BAY 11-7082 (3-[[4-(1,1-dimethylethyl)phenyl]-sulfonyl]-2-propenenitrile), BAY 11-7085 ((E)-(3-[[4-(1,1-dimethylethyl)phenyl]-sulfonyl]-2-propenenitrile), (see *Human Pathology*, 31(12), 1482 – 1490 (2000));
- cyclopentenone prostaglandin A1 (PGA1), (see *Proc. Natl. Acad. Sci. USA*, 94, 746-750 (1997)); and
- 10 - cyclopentabenzofurans as described in WO-00/08007.

Other NF- κ B inhibitors can be learned from literature and compounds can be tested for NF- κ B inhibiting activity using assays known in the art. See for instance *Proc. Natl. Acad. Sci. USA*, 94, 746-750 (1997).

The NF- κ B inhibitors for use in the treatment of mastitis can be specific inhibitors of NF- κ B or can be non-specific.

20 The present invention provides the use of a NF- κ B inhibitor for the manufacture of a medicament for the treatment of mastitis.

Consequently, the present invention provides a method for treating mastitis comprising administering to a subject an effective amount of a NF- κ B inhibitor.

25 As used herein, the term "subject" embraces warm-blooded animal species such as e.g. ruminants such as cattle, sheep, goats, buffalo etc.

There are a variety of forms or types of bovine mastitis, with varying severity and symptomatology, including the following :

- 30 - acute mastitis which is an abrupt and short-lived inflammation of the udder,
- chronic mastitis which is a moderate and persistent inflammation of the mammary gland.

35 Acute mastitis is generally a clinical form whereas chronic mastitis is often a subclinical form of the disease. Subclinical mastitis is a form of mastitis in which there is no swelling of the gland or observable abnormality of the milk, although there are

- changes in the milk, such as an increase in somatic cell count (SCC), that can be detected by special tests. This type of mastitis is by far the most prevalent and causes the greatest overall loss in most herds. It often is referred to as "hidden" mastitis. Clinical mastitis is a form of mastitis in which the abnormal conditions of the udder and secretion are observable. Mild clinical mastitis involves changes in the milk such as flakes, clots, and a watery or unusual appearance. Heat and sensitiveness of the udder are slight or absent, but there may be signs of swelling. Severe clinical mastitis involves a sudden onset with swelling of the infected quarter which is hot, hard and sensitive. The milk appears abnormal and milk production drops. Sometimes, in addition to the local effects in the udder, the cow herself becomes sick. There are signs of fever, rapid pulse, depression, weakness and loss of appetite. The combination of these conditions often is referred to as acute systemic mastitis, because not only the udder, but the whole animal is affected.
- 15 The effective amount of NF- κ B inhibitor for the treatment of mastitis will be determined using routine optimization techniques and are dependent upon the particular condition to be treated, the condition of the subject, the route of administration, the formulation, and the judgment of the practitioner and other factors evident to those skilled in the art. An effective amount may be achieved by multiple dosing.
- 20 Compositions comprising a NF- κ B inhibitor may be administered intravascularly, intraperitoneally, subcutaneously, intramuscularly, or by intramammary infusion using forms known in the pharmaceutical art. For intravascular, intraperitoneal, subcutaneous, intramuscular or intramammary administration, active drug components
- 25 may be combined with a suitable carrier such as water, saline, aqueous dextrose, and the like. Veterinary formulations comprising a NF- κ B inhibitor are usually prepared by mixing said NF- κ B inhibitor with an excipient, diluting by an excipient or enclosing within a carrier that can be in the form of a capsule, sachet, or other container. When the excipient serves as a diluent, it can be a solid, semi-solid, or liquid material that acts
- 30 as a vehicle, carrier or medium for the NF- κ B inhibitor. Examples of suitable excipients include lactose, dextrose, sucrose, sorbitol, mannitol, starches, gum acacia, calcium phosphate, alginates, tragacanth, gelatin, calcium silicate, microcrystalline cellulose, polyvinylpyrrolidone, cellulose, water, syrup, and methyl cellulose. The formulations can additionally include: lubricating agents such as talc, magnesium
- 35 stearate, and mineral oil; wetting agents; emulsifying and suspending agents; preserving agents such as methyl- and propylhydroxybenzoates; sweetening agents; and flavoring

agents. The compositions of the invention can be formulated to provide quick, sustained or delayed release of the active ingredient after administration to the host by procedures known in the art.

- 5 Intramammary infusions may be oil based, e.g. a vegetable oil such as peanut oil or a non-vegetable oil such as mineral oil and may further include a thickening agent and optionally also a surfactant.

10 Compositions can be in the form of tablets, pills, powders, lozenges, sachets, cachets, elixirs, suspensions, emulsions, solutions, infusions, syrups, aerosols, ointments, soft and hard gelatin capsules, suppositories, sterile injectable solutions and sterile packages powders. Regardless of the route of administration selected, these compositions are formulated into pharmaceutically acceptable dosage forms by conventional methods known to those skilled in the art.

15 The NF- κ B inhibitors may also be used in combination therapy with antibiotics. Antibiotics used in the treatment of mastitis are e.g. beta-lactam antibiotics such as ampicillin, cloxacillin, hetacillin, nafcillin, penicillin G, procaine penicillin; cephalosporins such as cefoperazone, cefuroxime, cefalonium, cefapirin, cefoxazole, 20 cefracetrile; aminoglycoside antibiotics such as framycetin, neomycin, novobiocin, streptomycin; macrolide antibiotics such as erythromycin; tetracyclines such as chlortetracycline, oxytetracycline; or polypeptide antibiotics such as polymyxin B.

Experimental details

- 25 The level of NF- κ B activity in milk cells from healthy and acute and chronic mastitis-affected cows was measured.

A. Measurement of NF- κ B activity in mastitis affected cows

1. Animals

- 30 Twenty healthy, six acute and twenty chronic mastitis-affected Holstein-Friesian cows were used in this study. Healthy cows had low somatic cell count (SCC) ($<10^5$ cells/ml) and were free of any clinical sign of mastitis. Acute mastitis-affected cows had high SCC ($> 4 \times 10^5$ cells/ml) and exhibited clinical signs of illness. Neither healthy nor acute mastitis-affected cows had earlier record of udder disease, as measured by clinical 35 symptoms and monthly SCC recordings. Chronic mastitis-affected cows had persistently increased SCC ($> 4 \times 10^5$ cells/ml), as determined by monthly SCC measurements, but were devoid of any clinical sign of the disease. All the cows were

between 2 and 7 years old, had a lactation number ranging from 1 to 5 and were lactating for 1 to 6 months. None of the cows received any treatment during the month preceding the experiments.

5 2. Sterile Milk Sampling

Milk was collected using a sterile teat cannula infusion apparatus. This apparatus consisted of a sterile and pyrogen-free cannula inserted into the teat canal and connected to the free end of an infusion set attached to a 2000 ml sterile collection bag. Milk samples were taken from one quarter. Before sampling, the quarter was
10 thoroughly disinfected with a solution containing 70% alcohol and 30% chlorohexidine. Immediately after milk sampling, an aliquot of 50 µl was the subject of a microbiological analysis. Samples that were found to be contaminated by bacteria were excluded from the study.

15 3. Somatic Cell Count

Immediately after milk collection, 50 ml of each milk sample was shipped to a specialized laboratory (Laboratory of the Milk Committee, Battice, Belgium), where a somatic cell count was performed following classical procedure.

20 4. Isolation of Milk Cells

Milk samples were first centrifuged (300 g) for 30 minutes at 4°C. Pellets were then washed in phosphate buffered saline (PBS) and centrifuged (300 g) for another 30 minutes at 4°C. The pelleted cells were finally resuspended in 1 ml of PBS before protein extraction. Cell differentials were performed on cytopsin preparations stained
25 with Diff-Quick (Dade Behring, Dudingon, Germany).

5. Protein Extraction

Isolated milk cells were centrifuged for 10 minutes at 300 g and the pellet was resuspended in 600 µl lysis buffer (0.1% nonidet P-40; 10 mM Tris HCl, pH 7.4;
30 10 mM NaCl; 3 mM MgCl₂; 1 mM EDTA; 2 mM orthovanadate; 1 mM diisopropyl fluorophosphates (DFP); 10 µg/ml leupeptin; 10 µg/ml aprotinin). The cells were vortexed 15 seconds, kept on ice for 5 minutes, and centrifuged for 10 minutes at 300 g. The resulting pellet was resuspended in a KCl buffer (10 mM Hepes, pH 7.4; 400 mM KCl; 10% Glycerol; 2 mM EDTA; 1 mM Ethylene glycol-bis[β-aminoethyl ether]-N,
35 N, N'-tetraacetic acid (EGTA); 1% nonidet P-40, 1 mM dithiothreitol (DTT); 2 mM orthovanadate; 10 µg/ml leupeptin; 10 µg/ml aprotinin; 1 mM DFP) and kept at 4°C

for 10 minutes before centrifugation for 15 minutes at 12,000 g. The supernatant was diluted three times and stored at -80°C until use.

6. NF-κB Electrophoretic Mobility Shift Assays (EMSAs)

5 Binding reactions were performed for 30 minutes at room temperature with 5 μg of total protein extracts in 20 mM Hepes (pH 7.9), 10 mM KCl, 0.2 mM EDTA, 20% (v/v) glycerol, 1% (w/v) acetylated bovine serum albumin, 3 μg of poly(dI-dC) (Amersham Pharmacia Biotech, Aylesbury, U.K.), 1 mM DTT (dithiothreitol), 1 mM phenylmethylsulfonyl fluoride, and 100,000 cpm of ³²P-labeled double-stranded
10 oligonucleotide probes. Probes were prepared by annealing the appropriate single-stranded oligonucleotides (Eurogentech, Liège, Belgium) at 65°C for 10 minutes in 10 mM Tris, 1 mM EDTA, and 10 mM NaCl, followed by slow cooling to room temperature. The probes were labeled by end-filling with the Klenow fragment of *E. coli* DNA polymerase I (Roche, Mannheim, Germany), with [³²P]dATP and
15 [³²P]dCTP (Dupont-New England Nuclear, Les Ulis, France). Labeled probes were purified by spin chromatography on Sephadex G-25 columns (Roche). DNA-protein complexes were separated from unbound probe on 4% native polyacrylamide gels at 150 V in 0.25 M Tris, 0.25 M sodium borate, and 0.5 mM EDTA, pH 8.0. Gels were vacuum-dried and exposed to x-ray film at -80°C for 12 hours. The amount of specific
20 complexes was determined by photodensitometry of the autoradiography. To confirm specificity, competition assays were performed with a 50-fold excess of unlabeled wild-type and mutated probes. For supershift experiments, 1.5 μl of each antibody was incubated with the extracts 30 minutes before addition of the radiolabeled probe. The sequences of the oligonucleotides used in this work were as follows : wild-type
25 palindromic κB probe, 5'-TTG GCA ACG GCA GGG GAA TTC CCC TCT CCT TAG GTT-3'; mutated palindromic κB probe, 5'-TTG GCA ACG GCA GAT CTA TTC CCC TCT CCT TAG GTT-3'.
The NF-κB activity was assessed by photodensitometry of the specific bands.

7. Results

30 Active NF-κB complexes were undetectable in milk cells from healthy cows, as determined by EMSAs. Conversely, protein extracts prepared from milk samples of acute mastitis-affected cows demonstrated a high NF-κB activity. In milk cells from cows suffering from chronic mastitis, NF-κB activity varied from low to high.
35

Linear regressions were carried out to assess the relation between the level of NF- κ B activity in milk cells, as assessed by photodensitometry, and the degree of inflammation in the mammary gland. This regression analysis demonstrated a highly significant ($P < 0.01$) correlation between NF- κ B activity and two markers of udder inflammation, namely SCC (Figure 1A) and the percentage of neutrophils in milk samples (Figure 1B).

This study demonstrated that NF- κ B activity was increased in milk cells from mastitis-affected cows and therefore NF- κ B inhibitors are believed to be of potential use in the treatment of mastitis.

B. Effects of administration of an NF- κ B inhibitor on milk cells obtained from chronic mastitis-affected cows

1. Animals

Four chronic mastitis-affected Holstein-Friesian cows were used in this study. These cows had persistently increased SCC ($> 4 \times 10^5$ cells/ml), as determined by monthly SCC measurements, but were devoid of any clinical sign of the disease. None of the cows received any treatment during the month preceding the experiments.

2. Sterile Milk Sampling

Milk was collected as described above.

3. Somatic Cell Count

Immediately after milk collection, 50 ml of each milk sample was shipped to a specialized laboratory (Laboratory of the Milk Committee, Battice, Belgium), where a somatic cell count was performed following classical procedure.

4. Isolation of Milk Cells

Milk samples were first centrifuged (300 g) for 30 minutes at 4°C. Pellets were then washed in phosphate buffered saline (PBS) and centrifuged (300 g) for another 30 minutes at 4°C. The pelleted cells were finally resuspended in 1 ml of PBS before protein extraction. Cell differentials were performed on cytopsin preparations stained with Diff-Quick (Dade Behring, Dudingen, Germany).

5. Culture and treatment of milk cells

Milk cells were cultured at 4×10^6 per ml in RPMI 1640 medium (Life Technologies, Merelbeke, Belgium) supplemented with 1% glutamine, 10% foetal bovine serum,

streptomycine 50 µg/ml, and penicillin 50 UI/ml. The cells were first incubated at 37°C in a 5% CO₂-95% air mixture for 90 minutes then treated with various doses of NF-κB inhibitors (15-dPGJ2 at concentrations of 40 µM, 50 µM, 60 µM and 80 µM; or gliotoxine at a concentration of 1µl/ml). Treated cells were incubated 3 hours before
5 protein extraction.

6. Protein Extraction

See procedure described above.

10 7. NF-κB Electrophoretic Mobility Shift Assays (EMSAs)

See procedure described above for assessing NF-κB activity by photodensitometry.

8. Results

15 In untreated milk cells from chronic mastitis-affected cows, an elevated level of NF-κB activity was detected, as determined by EMSAs. Conversely, protein extracts prepared from milk cells treated with 15d-PGJ2 or glyotoxine demonstrated a reduced NF-κB activity.

Description of the drawings

20 Figure 1 shows the relation between specific NF-κB activity displayed by milk cells, as determined by photodensitometry, and SCC (A) and the percentage of neutrophils (B) in healthy (■, n = 6) and acute (●, n = 6) and chronic (▲, n = 20) mastitis-affected Holstein-Friesian cows. r, correlation coefficient.

25 Figure 2 : 15d-PGJ2 and gliotoxine inhibit NF-κB activation in milk cells from chronic mastitis-affected cows. After isolation, milk cells were incubated for 90 min and then treated with the indicated concentrations of 15d-PGJ2 and gliotoxine. Protein extracts were assessed 3h after the treatment for NF-κB DNA-binding activity by EMSAs.

14. 08. 2002

92

CLAIMS

- 5 1. Use of a NF- κ B inhibitor for the manufacture of a medicament for the treatment of mastitis.
2. Use according to claim 1 wherein mastitis is chronic mastitis.
3. Use according to claim 1 wherein mastitis is acute mastitis.
- 10 4. Use according to any one of claims 1 to 3 in combination with antibiotics.
5. Use according to any one of the preceding claims wherein the NF- κ B inhibitor is a specific NF- κ B inhibitor .
- 15 6. Use according to any one of the preceding claims wherein the NF- κ B inhibitor is selected from 15d-PGJ2, tepoxalin, cycloepoxydon, (+)-cycloepoxydon, (-)-cycloepoxydon , sodium diethyldithiocarbamate, gliotoxin, MG-132, BAY 11-7082 or BAY 11-7085.
- 20 7. Use according to any one of the preceding claims wherein the NF- κ B inhibitor is administered intramammary.

ABSTRACT

92

5 USE OF NF-KAPPA B INHIBITORS FOR THE TREATMENT OF MASTITIS

The present invention relates to the use of NF- κ B (NF-kappa B) inhibitors for the manufacture of a medicament for the treatment of mastitis.

Figure 1 : Relation between specific NF- κ B activity displayed by milk cells, as determined by photodensitometry, and SCC (A) and the percentage of neutrophils (B) in healthy (■, $n = 6$) and acute (●, $n = 6$) and chronic (▲, $n = 20$) mastitis-affected Holstein-Friesian cows.
 r , correlation coefficient.




Figure 2 : 15d-PGJ2 and gliotoxine inhibit NF- κ B activation in milk cells from chronic mastitis-affected cows. After isolation, milk cells were incubated for 90 min and then treated with the indicated concentrations of 15d-PGJ2 and gliotoxine. Protein extracts were assessed 3h after the treatment for NF- κ B DNA-binding activity by EMSAs.

1-3-6 : protein extracts obtain from untreated milk neutrophils


2 : protein extracts obtain from milk neutrophils treated with 15-dPGJ2 40 μ M

4 : protein extracts obtain from milk neutrophils treated with 15-dPGJ2 50 μ M

5 : protein extracts obtain from milk neutrophils treated with 15-dPGJ2 60 μ M

7 : protein extracts obtain from milk neutrophils treated with gliotoxine 1 μ l/ml

8 : protein extracts obtain from milk neutrophils treated with 15-dPGJ2 80 μ M



**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☒ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☒ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.